



Mid-State Gastroenterology

powered by



PRE-PROCEDURE DAY INFORMATION

At least two days prior to your procedure, someone from the surgery center will call you before your procedure to confirm your arrival time and review information regarding your procedure.

PLEASE BE ADVISED OF THE FOLLOWING:

It is your responsibility to contact your insurance company to learn whether a specific procedure is covered and if it will be applied to a deductible. Some insurance plans have riders and underwriting on the plan that may or may not cover any or all of the procedure. We want you to be aware and understand the specifics of your insurance coverage. We are happy to provide you with the codes that could be used when submitting your claim to the insurance company.

*****VERY IMPORTANT*****

There are three (3) entities involved that will send a bill following your procedure.

1. FACILITY: Mid-State Endoscopy or Ascension Saint Thomas Rutherford
2. OneGI:
 - Dr. Sunil Sarvaria
 - Pathology (if any specimens are collected during procedure)
3. ANESTHESIA

Anesthesia, pathology, and physician charges are filed with insurance for determination of additional patient financial responsibility.

SURGERY CENTER LOCATIONS:

You will report to one of the surgery centers noted below as instructed by your provider/nurse.

**Mid-State Endoscopy
Center** 1115A Dow Street
Murfreesboro, TN 37130
Phone: 615-848-9234



Ascension Saint Thomas Rutherford
1700 Medical Center Pkwy
Murfreesboro, TN 37130
Phone: 615-396-4100

Enter through the main hospital entrance.



PROCEDURE DATE: _____

PROCEDURE TIME: _____

SuFlave Instructions

Medication has been electronically sent into your pharmacy, **PLEASE pick up within 5-7 days.**

PLEASE drink the required liquid to ensure a good clean out.

The **day prior** to your procedure, you will be on CLEAR LIQUIDS all day long.

DATE:

6PM the night before your procedure you will start your prep for your colonoscopy.

You will complete 1 bottle of 32 ounces of Suflave in one hour, you should drink 8 ounces every 15 minutes.

After completion of the Suflave bottle, you need to drink an additional 16 ounces of clear liquids.

Continue clear liquids till bedtime to ensure a good hydration.

6PM prior to your procedure you will start the second part of your prep.

DATE:

Complete the second bottle of 32 ounces of Suflave in one hour, you should drink 8 ounces every 15 minutes.

After completion of the Suflave bottle, you need to drink an additional 16 ounces of clear liquids.

NOTHING BY MOUTH FOUR HOURS PRIOR TO YOUR PROCEDURE, including gum, mints, candy, cough, drops, and water.

No usage of marijuana, cocaine, recreational drugs, or alcohol within 24 hours of your procedure. Anesthesia will not sedate you if you have consumed any of these in 24 hours.

PRODUCTS TO AVOID PRIOR TO YOUR PROCEDURE
(IMPORTANT: These are drugs that can thin your blood and may cause bleeding.)
Avoid these medications for at least 5 days before your procedure.

Advil	Loritab ASA
Aggrenox	Magsal
Aleve	Midol
Alka Seltzer	Mobic/Meloxicam
Anaprox	Mobigesic
Anacin	Monogesic Tablets
Arthritis Pain Formula	Motrin
Ascriptin	Naprosyn
Aspergum	Naprelan
Aspirin 325mg (Baby Aspirin 81mg is OK)	Naproxen
Azoid	Norgesic
BC Powders	Norwich
Bextra	Nuprin
Brilinta	P-A-C Analgesic
Bufferin	Pepto-Bismal
Carna Arthritis Medication	Percodan
Celebrex	Phentermine HCl
Clinoril	Plaquenil
Coumadin	Plavix
Darvon Compound /Darvocet	Pradaxa
Disalcid	Robaxisal
Doan's Pills	Salflex
Dristan	Salsalate
Easpirin	Salsitab
Ecotrin	Savaysa
Effervescent Tablets	Sine-Aid
Effient	Soma Compound
Eliquis	Stanback
Empirin	Synalgos-DC
Equagesic	Talwin
Etodolac	Therapy Bayer Caplets
Excedrin	Tolectin
Feldene	Tolmetin
Fenoprex	Toradol
Fiorinal	Trigesic
Fish Oil	Trilisate
Formula Caplets	Urisinus-Inlav
Goody Powders	Vanquish
Haltran	Vitamin E
Ibuprofen	Vioxx
Ibu-Tab	Warfarin
Indocin	Xarelto
Lodine	Zontivity

GLP-1 AGONISTS Type 2 diabetes or weight loss medications

Daily

- Rybeisus (semaglutide)
- Victoza (liraglutide)
- Saxenda (liraglutide)
- Byetta (exenatide)
- Adlyxin (Lixisenatide)

Weekly

- Ozempic (semaglutide)
- Wegovy (semaglutide)
- Trulicity (dulaglutide)
- Bydureon (exenatide)
- Mounjaro (tirzepatide)

Hold GLP-1 agonists on the **DAY** of procedure for patients who take the medication **DAILY**.

Hold GLP-1 agonists a **WEEK** prior to the procedure for the patients who take the medication **WEEKLY**.